




Contents lists available at SciVerse ScienceDirect

Molecular and Cellular Neuroscience

journal homepage: www.elsevier.com/locate/ymcne

Peripheral blood mono-nuclear cells derived from Alzheimer's disease patients show elevated baseline levels of secreted cytokines but resist stimulation with β -amyloid peptide

Natália Pessoa Rocha ^a, Antônio Lúcio Teixeira ^b, Fernanda Matos Coelho ^a, Paulo Caramelli ^b, Henrique Cerqueira Guimarães ^b, Izabela Guimarães Barbosa ^a, Tarcília Aparecida da Silva ^a, Marat Alexandrovich Mukhamedyarov ^c, Andrey Lvovich Zefirov ^c, Albert Anatolyevich Rizvanov ^{d,e,f,g}, Andrey Pavlovich Kiyasov ^{g,h}, Luciene B. Vieira ^a, Zoltán Janka ⁱ, András Palotás ^{j,*}, Helton José Reis ^a

^a Laboratório de Neurofarmacologia, Departamento de Farmacologia, Instituto de Ciências Biológicas, Universidade Federal de Minas Gerais (ICB-UFMG), Avenida Antônio Carlos 6627, 31270-901 Campus Pampulha, Belo Horizonte, Minas Gerais, Brazil

^b Departamento de Clínica Médica, Faculdade de Medicina, Universidade Federal de Minas Gerais, Avenida Alfredo Balena 190, 30130-100 Belo Horizonte, Minas Gerais, Brazil

^c Department of Physiology, Kazan State Medical University, ul. Butlerova 49, R-420012 Kazan, Russia

^d Department of Genetics, Faculty of Biology and Soil Sciences, Kazan Federal University, ul. Kremlevskaya 18, R-420008 Kazan, Russia

^e Core Research Laboratory, Kazan State Medical University, ul. Butlerova 49, R-420012 Kazan, Russia

^f Department of Genetics and Bioengineering, College of Engineering and Architecture, Yeditepe University, 26 Ağustos Campus, Kayisdagi cad., Kayisdagi, 34755 Istanbul, Turkey

^g Republic Clinical Hospital, ul. Orenburg tract 138, R-420064 Kazan, Russia

^h Department of Anatomy, Kazan State Medical University, ul. Butlerova 49, R-420012 Kazan, Russia

ⁱ Department of Psychiatry, Albert Szent-Györgyi Clinical Center, Faculty of Medicine, University of Szeged, H-6721 Szeged, Semmelweis u. 6, Hungary

^j Asklepios-Med (private medical practice and research center), H-6722 Szeged, Kossuth Lajos sgt. 23, Hungary

ARTICLE INFO

Article history:

Received 25 November 2010

Revised 11 September 2011

Accepted 19 September 2011

Available online 25 September 2011

Keywords:

Aging

Alzheimer's disease

β -amyloid peptide

Cytokine

Inflammaging

Peripheral blood mono-nuclear cell (PBMC)

ABSTRACT

Objectives: Among several other factors, the neuro-toxic β -amyloid peptide (β AP)-induced inflammatory mechanisms have also been implicated in the pathogenesis of Alzheimer's dementia (AD). Cytokines have recently emerged as prime candidates underlying this immune reaction. The purpose of this study was to evaluate the inflammatory response of peripheral blood mono-nuclear cells (PBMC) in AD.

Design: Cross-sectional (observational) study.

Setting: Behavioral and cognitive neurology clinic of the Universidade Federal de Minas Gerais in Belo Horizonte, Brazil.

Participants: AD patients ($n = 19$), healthy elderly ($n = 19$) and young ($n = 14$) individuals.

Measurements: Cytokine levels were assessed by enzyme-linked immuno-sorbent assay (ELISA) after exposing cells to a broad range of β AP concentrations (10^{-4} – 10^{-10} M) as a stimulus. AD samples were weighed against leukocytes harvested from non-demented young and elderly subjects.

Results: Cytokine production of PBMCs in the youth was characterized by low baseline levels when compared to cells from the older generation. In the aging population, AD cells were distinguished from the healthy elderly sub-group by an even higher basal cytokine secretion. The low resting concentration in young individuals was markedly increased after treatment with β AP, however cells from the elderly, irrespective of their disease status, showed unchanged cytokine release following β AP administration. Non-specific activation of PBMCs with anti-CD3/CD28 antibodies resulted in elevated interleukin (IL)-1 β concentrations in AD.

Conclusions: These results demonstrate a general over-production of cytokines and resistance to β AP in the old comparison group, with a more pronounced disruption/boosted pattern in AD. Our findings are in line with the hypothesis of "inflammaging", i.e. an enhanced inflammatory profile with normal aging and a further perturbed environment in AD. The observed cytokine profiles may serve as diagnostic biomarkers in dementia.

© 2011 Elsevier Inc. All rights reserved.

Introduction

Alzheimer's disease (AD) is a neuro-degenerative disorder characterized by progressive loss of cognitive functions, especially memory (Alzheimer's Association, 2009). The main histo-pathological picture consists of neuronal loss in selected brain regions, as well as deposition

* Corresponding author at: Asklepios-Med, H-6722 Szeged, Kossuth Lajos sgt. 23, Hungary.

E-mail address: palotas@asklepios-med.eu (A. Palotás).